

Diagnostic Imaging Pathways - Cholecystitis (Suspected Acute)

Population Covered By The Guidance

This pathway provides guidance for appropriate imaging investigation of adult patients with acute right upper quadrant pain or suspected acute cholecystitis.

Date reviewed: February 2015

Date of next review: 2017/2018

Published: July 2015

Quick User Guide

Move the mouse cursor over the **PINK** text boxes inside the flow chart to bring up a pop up box with salient points.

Clicking on the **PINK** text box will bring up the full text.

The relative radiation level (RRL) of each imaging investigation is displayed in the pop up box.

SYMBOL	RRL	EFFECTIVE DOSE RANGE
	None	0
	Minimal	< 1 millisieverts
	Low	1-5 mSv
	Medium	5-10 mSv
	High	>10 mSv

Pathway Diagram



What do I need to know?

1. Is the **diagnostic criteria** suggestive of acute cholecystitis?
2. Has the **severity assessment criteria** graded?
2. Is the patient elderly / critically ill / has multiple co-morbidities / nursed in ICU?
3. Is the patient possibly pregnant?

**CHOLECYSTITIS
 (SUSPECTED ACUTE) / RIGHT
 UPPER QUADRANT PAIN**

Date reviewed: February 2015
 Please note that this pathway is
 subject to review and revision




Image Gallery

Note: These images open in a new page

1 **Acute Cholecystitis**



Image 1 (Ultrasound): Grossly thickened gallbladder wall (12.8mm measured between callipers, normally up to 2-3mm) with calculi (arrow) within the gallbladder and pericholecystic fluid.

2 **Acute cholecystitis**



Image 2: Cholecystectomy showing acute cholecystitis with gallbladder wall oedema, vascular congestion and purulent exudate (blue arrow) caused by a massive cholesterol stone.

3 **Acute Cholecystitis**



Image 3 (H&E, x2.5): Histological section of severe acute cholecystitis showing extensive ulceration of the mucosa, haemorrhage, oedema and a dense transmural infiltrate of neutrophils and mononuclear inflammatory cells.

4 **Normal HIDA Scan**




Image 4 (HIDA Scan): Normal scan with visualisation of the gallbladder (GB) indicative of a patent cystic duct. The common bile duct (CBD) is also clearly identified.

Teaching Points

- Acute cholecystitis is the commonest cause of acute right upper quadrant (RUQ) pain, but other conditions such as peptic ulcer disease, pancreatitis and acute hepatitis can mimic acute cholecystitis
- Abdominal radiographs (AXR) in suspected acute cholecystitis are of no value [1](#)
- A chest radiograph may be performed to exclude a thoracic cause of pain
- Ultrasound is the preferred primary imaging test in suspected acute cholecystitis [2](#)
- Accuracy of CT in suspected acute cholecystitis is limited but literature has shown results were similar to ultrasound [3](#)
- Approximately 90-95% of acute cholecystitis is gallstones related, with acalculous disease accounting for 5-10% of cases. However CT is less sensitive than ultrasound in the detection of gallstones (57%-88%) [4](#)
- Indications for CT
 - If an alternative cause is considered
 - To identify complications
 - If clinical picture becomes non-specific
- The use of magnetic resonance imaging (MRI) is limited but likely, similar accuracy to ultrasound [2](#)
- Cholescintigraphy (Tc-IDA Radionuclide) scan is highly accurate but involves ionizing radiation, is a longer study and unreliable in severe hepatocellular disease. Used to clarify a negative, equivocal or technically difficult ultrasound if clinical suspicion of persists [2](#)

- Early laparoscopic cholecystectomy is the therapy of choice in operable patients [5](#)
- Image-guided percutaneous drainage of the gallbladder (percutaneous cholecystostomy), usually under ultrasound guidance has been extensively used for treating surgically unfit patients (elderly, critically ill and / or with severe comorbidities) with acute cholecystitis [6](#)
- Complications of acute cholecystitis include [7](#)
 - Gangrenous gallbladder
 - Gallbladder perforation
 - Pericholecystic abscess
 - Cholangitis
 - Fistula from gallbladder to bowel

Suspected Acute Cholecystitis / Right Upper Quadrant Pain

- Acute cholecystitis is the commonest cause of acute right upper quadrant pain (RUQ), but other conditions such as peptic ulcer disease, pancreatitis and acute hepatitis can mimic acute cholecystitis
- Diagnostic criteria have been defined by Japanese authors [8](#) recently as
 - A. Clinical: Local signs of inflammation, etc.

1. Murphy's sign
2. RUQ mass / pain / tenderness

B. Systemic signs of inflammation, etc.

1. fever
2. elevated CRP
3. abnormal WBC count

C. Imaging findings: imaging findings characteristic of acute cholecystitis

A definite diagnosis can be made in the presence of

1. one item in each A and B are positive
2. C confirms the diagnosis when acute cholecystitis is suspected clinically

Note: Acute hepatitis, other acute abdominal diseases, and chronic cholecystitis should be excluded

- Overall, in the setting of acute abdominal pain, CT has greater accuracy compared to ultrasonography. However in acute cholecystitis there is little data on the accuracy of CT; sensitivity and positive predictive value are probably similar [3](#)
- Ultrasound is the preferred primary imaging test in suspected acute cholecystitis as it has high accuracy, there is no ionizing radiation, it is more easily available (particularly at point-of-care), repeatable and images are not limited to the biliary tract (cf cholescintigraphy)
- A large multicenter randomized trial of early versus delayed laparoscopic cholecystectomy indicated that the former should become the therapy of choice for acute cholecystitis in operable patients [5](#)

Chest Radiograph

- The use of plain abdominal X-rays (AXR) in suspected acute cholecystitis are of no value [1](#)
- Most gallstones are non-radio-opaque. Conversely, visualization of opaque gallstones does not signify acute cholecystitis, since asymptomatic gallstones are common
- A chest radiograph may be performed to exclude a thoracic cause of pain [9](#)

Ultrasound

- Ultrasound is the preferred primary imaging test in suspected acute cholecystitis [10,11,12](#), although cholescintigraphy is more accurate [2](#)
- Published ultrasound studies show quite wide variation in accuracy, but a meta-analysis indicates a summary sensitivity and specificity for ultrasound are 82% and 81% respectively, compared to a sensitivity of 96% at a specificity of 90% for cholescintigraphy [2](#)
- However, ultrasound is favoured over cholescintigraphy as the primary imaging test as there is no ionizing radiation, it is more easily available (particularly at point-of-care), repeatable and images are not limited to the biliary tract, and is able to visualize gallstones with high sensitivity. It is also gives morphological information and is able to assess the bile ducts
- Ultrasound has been shown to be highly accurate for acute cholecystitis when the following signs are found in combination: gall bladder wall thickening / oedema, gall bladder distension, pericholecystic fluid, gallstones and positive Murphy sign. [12,13](#) However, recent studies in a regional hospital show less satisfactory accuracy [14,15](#)
- More than 90% diagnostic accuracy and varies with the morphologic criteria used [12,16,17,18](#)
- Colour / power Doppler increases accuracy over Gray-scale sonography [16](#)
- Advantages: allows evaluation of other abdominal structures (can identify an alternative diagnosis), provides preoperative information such as gallbladder size, stone size, gallbladder wall status and the presence of biliary dilatation [19](#)
- The addition of a HIDA scan in the diagnostic workup significantly improves sensitivity and can add valuable information in the appropriate clinical setting [15](#) particularly when ultrasound is negative or equivocal and there is continued clinical suspicion of acute cholecystitis
- In acute cholecystitis, the incidence of CBD stones was 1.8%. CBD diameter at ultrasound is not sufficient to identify patients with acute cholecystitis and CBD stones [20](#)

Treatment

- Ultrasound may predict the need for open versus laparoscopic cholecystectomy, typically in patients with severe acute cholecystitis found on ultrasound, combined with the following findings below, specifically when the duration of symptoms exceeds 3 days [21](#)
 - Gallbladder wall thickening to >5 mm
 - Pericholecystic exudates or abscess adjacent to the gallbladder
 - Difficulty identifying anatomical structures within Calot's triangle
- A large multicenter randomized trial of early versus delayed laparoscopic cholecystectomy indicated that the former should become the therapy of choice for acute cholecystitis (AC) in operable patients [5](#)

Percutaneous Cholecystostomy (PC)

- Image-guided percutaneous drainage of the gallbladder, usually under ultrasound guidance has been extensively used for treating surgically unfit patients with acute cholecystitis [6](#)
- Patients falling into this category are elderly, critically ill and / or with severe comorbidities. PC is a relatively safe procedure in these high-risk patients
- For calculous cholecystitis, however, there are no prospective trials of PC versus urgent cholecystectomy in these patients. A recent Cochrane database systematic review concluded that the role of PC could not be determined due to the lack of adequately powered randomized controlled trials. [22](#) A randomized controlled trial is currently underway of PC versus laparoscopic cholecystectomy in acute cholecystitis in high risk patients (the "CHOCOLATE" trial); results are awaited
- Of those patients who proceed to delayed cholecystectomy following PC, there is a high rate of conversion from laparoscopic to open cholecystectomy. [23](#) However, PC may be definitive treatment with delayed cholecystectomy being unnecessary [24,25,26](#)

Suspected Complications of Cholecystitis

- Patients with acute cholecystitis may develop complications such as gangrenous, perforated or emphysematous cholecystitis
- Worrisome ultrasound findings for complicated cholecystitis include the following below, where CT may add useful information in such patients [7](#)
 - Intraluminal findings (sloughed mucosa, haemorrhage and / or abnormal gas)
 - Gallbladder wall abnormalities (striations, asymmetrical wall thickening, abnormal gas, loss of sonorectivity and / or contrast enhancement)
 - Pericholecystic changes (echogenic fat, pericholecystic fluid and / or abscess formation)

Computed Tomography (CT)

- Data on the accuracy of CT in suspected acute cholecystitis is limited but in one comparative study versus ultrasound, the results were similar [3](#)
- CT findings of acute cholecystitis include gallbladder wall thickening, gallbladder distention, pericholecystic fat stranding, subserosal oedema, mucosal enhancement, transient focal enhancement of the liver adjacent to the gallbladder and pericholecystic fluid collection [27,28,29](#)
- Complicated acute cholecystitis can be diagnosed, including pericholecystic abscess and gas collection within gallbladder [28](#)
- Approximately 90-95% of acute cholecystitis is gallstones related, with acalculous disease accounting for 5-10% of cases
- CT is less sensitive than ultrasound in the detection of gallstones (57%-88%) [4](#)
- CT in acute right upper quadrant (RUQ) pain is best used as a secondary test if
 - The diagnosis of acute cholecystitis is unconfirmed or equivocal on initial US (\pm cholescintigraphy), and an alternative diagnosis is being considered
 - If complications such as gangrenous gallbladder, perforation and pericholecystic abscess or fistula formation are suspected
 - When the clinical picture is non-specific, as it can detect other intra-abdominal inflammatory processes and when complications of acute cholecystitis are suspected

Cholescintigraphy (Tc-IDA Radionuclide Scan)

- Highly accurate in the diagnosis of acute cholecystitis [2,17,18](#)
- Used to clarify a negative, equivocal or technically difficult ultrasound in the presence of continued clinical suspicion of acute cholecystitis [19](#)
- The hallmark of acute cholecystitis (calculous as well as acalculous) is persistent gall bladder non-visualisation 30 minutes post-morphine or on the 3-4 hour delayed image [19](#)
- False positives can occur in [30](#)
 - Alcoholics
 - Intensive care unit (ICU) patients
 - Patients fasting for a prolonged period
 - Cystic fibrosis
 - Chronic cholecystitis
- Morphine augmentation reduces false positives and is superior to delayed imaging [31,32,33](#)
- The drawbacks of cholescintigraphy are [19](#)
 - Ionizing radiation
 - Longer examination time (than ultrasound or CT)
 - Unreliable in severe hepatocellular disease or at serum bilirubin levels >340-500 mmol/L
 - Inability to diagnose extra-biliary causes of acute right upper quadrant abdominal pain
 - Inability to provide anatomical information

Magnetic Resonance Imaging (MRI)

There have been few studies of MRI in suspected acute cholecystitis. Those available suggest accuracy is similar to ultrasound. [2](#) However, the role of MRI in the acute setting is limited

References

Date of literature search: February 2015

The search methodology is available on request. [Email](#)

References are graded from Level I to V according to the Oxford Centre for Evidence-Based Medicine, Levels of Evidence. [Download the document](#)

- [1.](#) Bhangu A, Richardson C, Winter H, Bleetman A. **Value of initial radiological investigations in patients admitted to hospital with appendicitis, acute gallbladder disease or acute pancreatitis.** Emerg Med J. 2010 Oct;27(10):754-7. (Level III evidence). [View the reference](#)
- [2.](#) Kiewiet JJ, Leeuwenburgh MM, Bipat S, Bossuyt PM, Stoker J, Boermeester MA. **A systematic review and meta-analysis of diagnostic performance of imaging in acute cholecystitis.** Radiology. 2012;264(3):708-20. (Level II evidence). [View the reference](#)
- [3.](#) van Randen A, Lameris W, van Es HW, van Heesewijk HP, van Ramshorst B, Ten Hove W, et al. **A comparison of the accuracy of ultrasound and computed tomography in common diagnoses causing acute abdominal pain.** Eur Radiol. 2011;21(7):1535-45. (Level II evidence). [View the reference](#)
- [4.](#) Van Beers BE, Pringot JH. **Imaging of cholelithiasis: helical CT.** Abdom Imaging. 2001;26(1):15-20. (Review article). [View the reference](#)
- [5.](#) Gutt CN, Encke J, Koninger J, Harnoss JC, Weigand K, Kipfmuller K, et al. **Acute cholecystitis: early versus delayed cholecystectomy, a multicenter randomized trial (ACDC study, NCT00447304).** Ann Surg. 2013;258(3):385-93. (Level II evidence). [View the reference](#)
- [6.](#) Tsuyuguchi T, Itoi T, Takada T, Strasberg SM, Pitt HA, Kim MH, et al. **TG13 indications and**

- techniques for gallbladder drainage in acute cholecystitis (with videos).** J Hepatobiliary Pancreat Sci. 2013;20(1):81-8. (Guideline). [View the reference](#)
7. Charalel RA, Jeffrey RB, Shin LK. **Complicated cholecystitis: the complementary roles of sonography and computed tomography.** Ultrasound Q. 2011;27(3):161-70. (Review article). [View the reference](#)
8. Yokoe M, Takada T, Strasberg SM, Solomkin JS, Mayumi T, Gomi H, et al. **New diagnostic criteria and severity assessment of acute cholecystitis in revised Tokyo Guidelines.** J Hepatobiliary Pancreat Sci. 2012;19(5):578-85. (Guideline). [View the reference](#)
9. Field S, Guy PJ, Upsdell SM, Scourfield AE. **The erect abdominal radiograph in the acute abdomen: should its routine use be abandoned?** Br Med J (Clin Res Ed). 1985;290(6486):1934-6. (Level II evidence). [View the reference](#)
10. Laing FC, Federle MP, Jeffrey RB, Brown TW. **Ultrasonic evaluation of patients with acute right upper quadrant pain.** Radiology. 1981;140(2):449-55. (Level III evidence). [View the reference](#)
11. Harvey RT, Miller WT, Jr. **Acute biliary disease: initial CT and follow-up US versus initial US and follow-up CT.** Radiology. 1999;213(3):831-6. (Level III evidence). [View the reference](#)
12. Ralls PW, Colletti PM, Lapin SA, Chandrasoma P, Boswell WD, Jr., Ngo C, et al. **Real-time sonography in suspected acute cholecystitis. Prospective evaluation of primary and secondary signs.** Radiology. 1985;155(3):767-71. (Level II evidence). [View the reference](#)
13. Yarmenitis S. **Ultrasound of the gallbladder and the biliary tree.** Eur Radiol. 2002;12(2):270-82. (Review article). [View the reference](#)
14. Hwang H, Marsh I, Doyle J. **Does ultrasonography accurately diagnose acute cholecystitis? Improving diagnostic accuracy based on a review at a regional hospital.** Can J Surg. 2014;57(3):162-8. (Level II evidence). [View the reference](#)
15. Kaoutzanis C, Davies E, Leichtle SW, Welch KB, Winter S, Lampman RM, et al. **Abdominal ultrasound versus hepato-imino diacetic acid scan in diagnosing acute cholecystitis--what is the real benefit?** J Surg Res. 2014;188(1):44-52. (Level IV evidence). [View the reference](#)
16. Soyer P, Brouland JP, Boudiaf M, Kardache M, Pelage JP, Panis Y, et al. **Color velocity imaging and power Doppler sonography of the gallbladder wall: a new look at sonographic diagnosis of acute cholecystitis.** AJR Am J Roentgenol. 1998;171(1):183-8. (Level II/III evidence). [View the reference](#)
17. Chatziioannou SN, Moore WH, Ford PV, Dhekne RD. **Hepatobiliary scintigraphy is superior to abdominal ultrasonography in suspected acute cholecystitis.** Surgery. 2000;127(6):609-13. (Level II evidence). [View the reference](#)
18. Shea JA, Berlin JA, Escarce JJ, Clarke JR, Kinosian BP, Cabana MD, et al. **Revised estimates of diagnostic test sensitivity and specificity in suspected biliary tract disease.** Arch Intern Med. 1994;154(22):2573-81. (Level II/III evidence). [View the reference](#)
19. Lin EC, Kuni CC. **Radionuclide imaging of hepatic and biliary disease.** Semin Liver Dis. 2001;21(2):179-94. (Review article). [View the reference](#)
20. Boys JA, Doorly MG, Zehetner J, Dhanireddy KK, Senagore AJ. **Can ultrasound common bile duct diameter predict common bile duct stones in the setting of acute cholecystitis?** Am J Surg. 2014;207(3):432-5. (Level III evidence). [View the reference](#)
21. Cwik G, Skoczylas T, Wyrosiak-Najs J, Wallner G. **The value of percutaneous ultrasound in predicting conversion from laparoscopic to open cholecystectomy due to acute cholecystitis.** Surg Endosc. 2013;27(7):2561-8. (Level IV evidence). [View the reference](#)
22. Gurusamy KS, Davidson C, Gluud C, Davidson BR. **Early versus delayed laparoscopic cholecystectomy for people with acute cholecystitis.** Cochrane Database Syst Rev. 2013;6:Cd005440. (Review article). [View the reference](#)
23. Sanjay P, Mittapalli D, Marioud A, White RD, Ram R, Alijani A. **Clinical outcomes of a percutaneous cholecystostomy for acute cholecystitis: a multicentre analysis.** HPB (Oxford). 2013;15(7):511-6. (Level III evidence). [View the reference](#)
24. Li M, Li N, Ji W, Quan Z, Wan X, Wu X, et al. **Percutaneous cholecystostomy is a definitive**



treatment for acute cholecystitis in elderly high-risk patients. Am Surg. 2013;79(5):524-7. (Level IV evidence). [View the reference](#)

25. Cha BH, Song HH, Kim YN, Jeon WJ, Lee SJ, Kim JD, et al. **Percutaneous cholecystostomy is appropriate as definitive treatment for acute cholecystitis in critically ill patients: a single center, cross-sectional study.** Korean J Gastroenterol. 2014;63(1):32-8. (Level III evidence). [View the reference](#)

26. Chang YR, Ahn YJ, Jang JY, Kang MJ, Kwon W, Jung WH, et al. **Percutaneous cholecystostomy for acute cholecystitis in patients with high comorbidity and re-evaluation of treatment efficacy.** Surgery. 2014;155(4):615-22. (Level IV evidence). [View the reference](#)

27. Paulson EK. **Acute cholecystitis: CT findings.** Semin Ultrasound CT MR. 2000;21(1):56-63. (Review article). [View the reference](#)

28. Yokoe M, Takada T, Strasberg SM, Solomkin JS, Mayumi T, Gomi H, et al. **TG13 diagnostic criteria and severity grading of acute cholecystitis (with videos).** J Hepatobiliary Pancreat Sci. 2013;20(1):35-46. (Guideline). [View the reference](#)

29. Fidler J, Paulson EK, Layfield L. **CT evaluation of acute cholecystitis: findings and usefulness in diagnosis.** AJR Am J Roentgenol. 1996;166(5):1085-8. (Level III evidence). [View the reference](#)

30. Fig LM, Wahl RL, Stewart RE, Shapiro B. **Morphine-augmented hepatobiliary scintigraphy in the severely ill: caution is in order.** Radiology. 1990;175(2):467-73. (Level III evidence). [View the reference](#)

31. Flancbaum L, Choban PS, Sinha R, Jonasson O. **Morphine cholescintigraphy in the evaluation of hospitalized patients with suspected acute cholecystitis.** Ann Surg. 1994;220(1):25-31. (Level III evidence). [View the reference](#)

32. Flancbaum L, Choban PS. **Use of morphine cholescintigraphy in the diagnosis of acute cholecystitis in critically ill patients.** Intensive Care Med. 1995;21(2):120-4. (Level III evidence). [View the reference](#)

33. Kim CK, Tse KK, Juweid M, Mozley PD, Woda A, Alavi A. **Cholescintigraphy in the diagnosis of acute cholecystitis: morphine augmentation is superior to delayed imaging.** J Nucl Med. 1993;34(11):1866-70. (Level II evidence). [View the reference](#)

Information for Consumers

Information from this website	Information from the Royal Australian and New Zealand College of Radiologists' website
<p>Radiation Risks of X-rays and Scans</p> <p>Computed Tomography (CT)</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Ultrasound</p> <p>Chest Radiograph (X-ray)</p> <p>Plain Radiography (X-ray)</p>	<p>Computed Tomography (CT)</p> <p>Contrast Medium (Gadolinium versus Iodine)</p> <p>Gadolinium Contrast Medium</p> <p>Iodine-Containing Contrast Medium</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Plain Radiography/X-rays</p>



[Radiation Risk of Medical Imaging During Pregnancy](#)

[Radiation Risk of Medical Imaging for Adults and Children](#)

[Ultrasound](#)

[Nuclear Medicine](#)

Copyright

© Copyright 2015, Department of Health Western Australia. All Rights Reserved. This web site and its content has been prepared by The Department of Health, Western Australia. The information contained on this web site is protected by copyright.

Legal Notice

Please remember that this leaflet is intended as general information only. It is not definitive and The Department of Health, Western Australia can not accept any legal liability arising from its use. The information is kept as up to date and accurate as possible, but please be warned that it is always subject to change.

File Formats

Some documents for download on this website are in a Portable Document Format (PDF). To read these files you might need to download Adobe Acrobat Reader.



[Legal Matters](#)