



Diagnostic Imaging Pathways - Cholecystitis (Acalculous, Suspected Acute)

Population Covered By The Guidance

This pathway provides guidance for appropriate imaging investigation of critically ill patients in the Intensive Care Unit (ICU) setting presenting with right upper quadrant pain or when acute acalculous cholecystitis is suspected.

Date reviewed: February 2015

Date of next review: 2017/2018






Published: July 2015

Quick User Guide

Move the mouse cursor over the **PINK** text boxes inside the flow chart to bring up a pop up box with salient points.

Clicking on the **PINK** text box will bring up the full text.

The relative radiation level (RRL) of each imaging investigation is displayed in the pop up box.

SYMBOL	RRL	EFFECTIVE DOSE RANGE
	None	0
	Minimal	< 1 millisieverts
	Low	1-5 mSv
	Medium	5-10 mSv
	High	>10 mSv

Pathway Diagram




Image Gallery

This page is blank

Teaching Points

- Ultrasound is of limited use since in intensive care unit patients it has low specificity [1](#); gallbladder abnormalities are frequently seen on ultrasound in intensive care patients who do not have suspected AAC. [2](#) This is also true of CT. [3](#) Cholescintigraphy is also associated with false negative and false positive results in suspected AAC [4](#)
- Percutaneous cholecystostomy (PC) and drainage can be both diagnostic and therapeutic in AAC [5](#)

Suspected Acute Acalculous Cholecystitis (AAC)

- AAC is a complication of severe morbidity when primary diagnosis is not acute cholecystitis
- This is a serious condition that typically occurs in critically ill patients with severe comorbidities, usually in the intensive care environment. In this setting, AAC is difficult to diagnose

Percutaneous Cholecystostomy (PC)

- Percutaneous cholecystostomy (PC) and drainage can be both diagnostic and therapeutic in AAC. Essentially, a successful 'trial of therapy' by PC is presumptive of the diagnosis of AAC. A



subsequent delayed cholecystectomy may not be required

References

Date of literature search: February 2015

The search methodology is available on request. [Email](#)

References are graded from Level I to V according to the Oxford Centre for Evidence-Based Medicine, Levels of Evidence. [Download the document](#)

1. Puc MM, Tran HS, Wry PW, Ross SE. **Ultrasound is not a useful screening tool for acute acalculous cholecystitis in critically ill trauma patients.** Am Surg. 2002;68(1):65-9. (Level III evidence). [View the reference](#)
2. Boland GW, Slater G, Lu DS, Eisenberg P, Lee MJ, Mueller PR. **Prevalence and significance of gallbladder abnormalities seen on sonography in intensive care unit patients.** AJR Am J Roentgenol. 2000;174(4):973-7. (Level III evidence). [View the reference](#)
3. Ahvenjarvi L, Koivukangas V, Jartti A, Ohtonen P, Saarnio J, Syrjala H, et al. **Diagnostic accuracy of computed tomography imaging of surgically treated acute acalculous cholecystitis in critically ill patients.** J Trauma. 2011;70(1):183-8. (Level III evidence). [View the reference](#)
4. Ziessman HA. **Nuclear medicine hepatobiliary imaging.** Clin Gastroenterol Hepatol. 2010;8(2):111-6. (Review article). [View the reference](#)
5. Chung YH, Choi ER, Kim KM, Kim MJ, Lee JK, Lee KT, et al. **Can percutaneous cholecystostomy be a definitive management for acute acalculous cholecystitis?** J Clin Gastroenterol. 2012;46(3):216-9. (Level IV evidence). [View the reference](#)

Information for Consumers

Information from this website	Information from the Royal Australian and New Zealand College of Radiologists' website
<p>Radiation Risks of X-rays and Scans</p> <p>Computed Tomography (CT)</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Ultrasound</p> <p>Chest Radiograph (X-ray)</p> <p>Plain Radiography (X-ray)</p>	<p>Computed Tomography (CT)</p> <p>Contrast Medium (Gadolinium versus Iodine)</p> <p>Gadolinium Contrast Medium</p> <p>Iodine-Containing Contrast Medium</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Plain Radiography/X-rays</p> <p>Radiation Risk of Medical Imaging During</p>



	<p>Pregnancy</p> <p>Radiation Risk of Medical Imaging for Adults and Children</p> <p>Ultrasound</p> <p>Nuclear Medicine</p>
--	---

Copyright

© Copyright 2015, Department of Health Western Australia. All Rights Reserved. This web site and its content has been prepared by The Department of Health, Western Australia. The information contained on this web site is protected by copyright.

Legal Notice

Please remember that this leaflet is intended as general information only. It is not definitive and The Department of Health, Western Australia can not accept any legal liability arising from its use. The information is kept as up to date and accurate as possible, but please be warned that it is always subject to change

File Formats

Some documents for download on this website are in a Portable Document Format (PDF). To read these files you might need to download Adobe Acrobat Reader.



[Legal Matters](#)