

Diagnostic Imaging Pathways - Paediatric, Elbow Injury

Population Covered By The Guidance

This pathway provides guidance on imaging children with elbow injuries.

Date reviewed: July 2014

Date of next review: 2017/2018

Published: July 2014

Quick User Guide

Move the mouse cursor over the **PINK** text boxes inside the flow chart to bring up a pop up box with salient points.

Clicking on the **PINK** text box will bring up the full text.

The relative radiation level (RRL) of each imaging investigation is displayed in the pop up box.

SYMBOL	RRL	EFFECTIVE DOSE RANGE
	None	0
	Minimal	< 1 millisieverts
	Low	1-5 mSv
	Medium	5-10 mSv
	High	>10 mSv

Pathway Diagram

Date reviewed: July 2014
Please note that this pathway is subject to review and revision




Image Gallery

Note: Images coming soon

Teaching Points

- Trauma to the paediatric elbow is common, and there is a higher incidence in this population compared with adults

- Children instinctively protect themselves by using outstretched hands when they fall. 75% of all paediatric fractures occur in the upper extremity [1](#)
- Unrecognised fractures may result in permanent neurovascular deficit and growth disturbances
- Radiographic imaging is commonly used to assess elbow injuries

Plain Radiographs

- Plain radiography of the elbow is the investigative mainstay for assessing elbow injuries in children in the acute and sub-acute setting
- Interpretation is complicated by the relative radiolucency of the cartilaginous anatomy and complex nature of epiphyseal growth. Although ossification centres generally develop in sequence, there is variability with gender and race
- Standard views are antero-posterior & lateral projections of the elbow. Additional oblique views and comparison views of the opposite limb may also aid assessment. Stress views using valgus-stress can provide additional information on the ligamentous integrity of the joint in the sub-acute and chronic setting [2](#)
- In many cases, particularly those involving high-energy trauma, fractures may be readily apparent
- When no fracture is apparent, radiographic findings such as the presence of a joint effusion or displacement of the fat pads can raise the suspicion of an occult fracture
- Occult fractures are found in 80% of patients with isolated posterior fat pad displacement on initial or follow-up radiographs [3](#)
- The sensitivity and specificity of plain radiographs for detecting elbow fractures has been documented as 64-71% and 56% respectively [4,5](#)
- Whilst MRI scans detect more injuries than radiographs, the vast majority of these will make no difference to patient management or outcome

Computed Tomography (CT)

- Multidetector computed tomography (MDCT) allows rapid scanning times, multiplanar and 3D reconstruction and, unlike MRI, is readily available and generally does not require sedation or general anaesthesia
- Due to its excellent resolution of bony structures, MDCT may be used for fracture characterisation and for pre-operative planning of complex fractures
- A major limitation is the radiation dose which is of particular concern in the paediatric population. However, dose reduction techniques and low-dose protocols can substantially reduce the radiation exposure [6](#)
- Chapman et al found that MDCT was highly sensitive and specific (92%, 79% respectively) for detecting radiographically occult fractures of paediatric elbows [7](#)

Magnetic Resonance Imaging (MRI)

- Magnetic resonance imaging has the advantages of no ionising radiation, excellent soft tissue resolution and capable of multi planar reconstructions
- Rarely would an MRI scan be required. Whilst MRI may detect additional injuries these rarely change management
- MRI is able to evaluate a spectrum of injuries not visible on plain radiographs. These include injuries to the soft tissues, cartilage/epiphyses and bone contusions

- Disadvantages include costs, limited access and the need for sedation or general anaesthesia for young children
- There have been a number of case series examining the use of MRI in paediatric patients with elbow trauma. In general, MRI has been used when plain radiographs are equivocal or demonstrate an effusion only
 - Griffith et al studied 50 paediatric patients with moderately severe elbow trauma not requiring immediate surgery. They found radiographically occult fractures in 22% and radiographically occult transphyseal or physeal injury in 19%. However MRI findings did not result in any change in treatment [4](#)
 - Wei et al studied children with little league elbow and, whilst MRI demonstrated more abnormalities compared with radiographs, these did not change clinical management [8](#)
 - Pudas et al studied 25 consecutive paediatric patients with elbow trauma. In those patients with a radiographic joint effusion but no fracture, MRI confirmed occult fracture in 8 of 9 patients and led to a change in management of 4 of the 9 patients. Three of these involved putting the patient in a cast although one had septic arthritis subsequently diagnosed after a fracture was excluded [5](#)
 - Major et al studied 13 consecutive paediatric & adult patients with elbow trauma & radiographic joint effusion. 4 of the 7 children demonstrated occult fractures. All patients demonstrated bone marrow oedema [9](#)
 - Carey et al studied 14 patients with documented or suspected growth plate fractures of the upper and lower limbs. The MRI findings changed the diagnosis in 7 of 14 patients and management in 5 patients [10](#)
- MR and CT Arthrography are advanced joint imaging studies where contrast is injected directly into the joint space to study the internal architecture of the joint. These are reserved for conditions like capsular and ligamentous injuries, intra-articular loose bodies, focal cartilaginous abnormalities and elbow dislocations. These conditions tend to be associated with young athletes [12](#)

References

- [1.](#) Jacoby SM, Herman MJ, Morrison WB, Osterman AL. **Pediatric elbow trauma: an orthopaedic perspective on the importance of radiographic interpretation.** Semin Musculoskelet Radiol. 2007;11(1):48-56. (Review article)
- [2.](#) Potter HG, Ho ST, Altchek DW. **Magnetic resonance imaging of the elbow.** Semin Musculoskelet Radiol. 2004; 8(1):5-16. (Review article)
- [3.](#) Skaggs DL, Mirzayan R. **The posterior fat pad sign in association with occult fracture of the elbow in children.** J Bone Joint Surg Am. 1999;81(10):1429–33. (Level II/III evidence)
- [4.](#) Griffith JF, Roebuck DJ, Cheng JC, Chan YL, Rainer TH, Ng BK, Metreweli C. **Acute elbow trauma in children: spectrum of injury revealed by MR imaging not apparent on radiographs.** AJR Am J Roentgenol. 2001;176(1):53-60. (Level III evidence)
- [5.](#) Pudas T, Hurme T, Mattila K, Svedström E. **Magnetic resonance imaging in pediatric elbow fractures.** Acta Radiol. 2005;46(6):636-44. (Level III evidence)
- [6.](#) Chapman VM, Kalra M, Halpern E, Grottkau B, Albright M, Jaramillo D. **16-MDCT of the posttraumatic pediatric elbow: optimum parameters and associated radiation dose.** AJR Am J Roentgenol. 2005;185(2):516-21. (Level III evidence)
- [7.](#) Chapman V, Grottkau B, Albright M, Elaini A, Halpern E, Jaramillo D. **MDCT of the elbow in pediatric patients with posttraumatic elbow effusions.** AJR Am J Roentgenol. 2006;187(3):812-7. (Level III evidence)
- [8.](#) Wei A, Khana S, Limpisvasti O, Crues J, Podesta L, Yocum LA. **Clinical and magnetic resonance imaging findings associated with little league elbow.** J Pediatr Orthop. 2010;30(7):715-9. (Level II/III evidence)



9. Major NM, Crawford ST. **Elbow effusions in trauma in adults and children: is there an occult fracture?** AJR Am J Roentgenol. 2002;178(2):413-8. (Level II evidence)
10. Carey J, Spence L, Blickman H, Eustace S. **MRI of pediatric growth plate injury: correlation with plain film radiographs and clinical outcome.** Skeletal Radiol. 1998;27(5):250-5. (Level III evidence)
11. Iyer RS, Thapa MM, Khanna PC, Chew FS. **Paediatric bone imaging: imaging elbow trauma in children - a review of acute and chronic injuries.** AJR Am J Roentgenol. 2012; 198:1053-68. (Review article)
12. Delpont AG, Zoga AC. **MR and CT arthrography of the elbow.** Semin Musculoskelet Radiol. 2012;16:15–26. (Review article).

Information for Consumers

Information from this website	Information from the Royal Australian and New Zealand College of Radiologists' website
<p>Consent to Procedure or Treatment</p> <p>Radiation Risks of X-rays and Scans</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Plain Radiography (X-ray)</p>	<p>Magnetic Resonance Imaging (MRI)</p> <p>Plain Radiography/X-rays</p> <p>Radiation Risk of Medical Imaging for Adults and Children</p> <p>Children's (Paediatric) X-ray Examination</p> <p>Making Your Child's Test or Procedure Less Stressful</p>

Copyright

© Copyright 2015, Department of Health Western Australia. All Rights Reserved. This web site and its content has been prepared by The Department of Health, Western Australia. The information contained on this web site is protected by copyright.

Legal Notice

Please remember that this leaflet is intended as general information only. It is not definitive and The Department of Health, Western Australia can not accept any legal liability arising from its use. The information is kept as up to date and accurate as possible, but please be warned that it is always subject to change.



File Formats

Some documents for download on this website are in a Portable Document Format (PDF). To read these files you might need to download Adobe Acrobat Reader.



[Legal Matters](#)